

Maximal monotone operators are selfdual vector fields and vice-versa

Nassif Ghoussoub

Department of Mathematics, University of British Columbia,
Vancouver BC Canada V6T 1Z2
nassif@math.ubc.ca

February 2, 2008

Abstract

If L is a selfdual Lagrangian L on a reflexive phase space $X \times X^*$, then the vector field $x \rightarrow \bar{\partial}L(x) := \{p \in X^*; (p, x) \in \partial L(x, p)\}$ is maximal monotone. Conversely, any maximal monotone operator T on X is derived from such a potential on phase space, that is there exists a selfdual Lagrangian L on $X \times X^*$ (i.e., $L^*(p, x) = L(x, p)$) such that $T = \bar{\partial}L$.

This solution to problems raised by Fitzpatrick can be seen as an extension of a celebrated result of Rockafellar stating that maximal cyclically monotone operators are actually of the form $T = \partial\varphi$ for some convex lower semi-continuous function on X . This representation allows for the application of the selfdual variational theory –recently developed by the author– to the equations driven by maximal monotone vector fields. Consequently, solutions to equations of the form $\Lambda x \in Tx$ for a given map $\Lambda : D(\Lambda) \subset X \rightarrow X^*$, can now be obtained by minimizing functionals of the form $I(x) = L(x, \Lambda x) - \langle x, \Lambda x \rangle$.

1 Introduction

In a series of papers ([4]–[10]), we developed and used the concept of selfdual Lagrangians L on phase space $X \times X^*$ –where X is a Banach space– in order to provide variational formulations and resolutions to various differential equations which are not variational in the sense of Euler-Lagrange. The main idea behind this theory –which is developed in its full generality in the upcoming monograph [6]– originated from the fact that a large set of PDEs and evolution equations can be written in the form

$$(p, x) \in \partial L(x, p), \quad (1)$$

where ∂L is the subdifferential of a Lagrangian $L : X \times X^* \rightarrow \mathbb{R} \cup \{+\infty\}$ that is convex and lower semi-continuous –in both variables– while satisfying

$$L^*(p, x) = L(x, p) \text{ for all } (p, x) \in X^* \times X. \quad (2)$$

Here L^* is the Legendre transform of L in both variables, that is

$$L^*(p, x) = \sup \{ \langle p, y \rangle + \langle x, q \rangle - L(y, q); (y, q) \in X \times X^* \}.$$

Solutions are then found for a given p by simply minimizing the functional $I_p(x) = L(x, p) - \langle x, p \rangle$ and by proving that the minimum is actually zero. In other words, by defining the *derived vector fields of L* at $x \in X$ to be the –possibly empty– sets

$$\bar{\partial}L(x) := \{p \in X^*; L(x, p) - \langle x, p \rangle = 0\} = \{p \in X^*; (p, x) \in \partial L(x, p)\}, \quad (3)$$

one can then find variationally the zeroes of those set-valued maps $T : X \rightarrow 2^{X^*}$ of the form $T(x) = \bar{\partial}L(x)$ for some selfdual Lagrangian L on $X \times X^*$.

These *selfdual vector fields* are natural extensions of subdifferentials of convex lower semi-continuous functions. Indeed, the most basic selfdual Lagrangians are of the form $L(x, p) = \varphi(x) + \varphi^*(p)$ where φ is such a function on X , and φ^* is its Legendre conjugate on X^* , in which case $\bar{\partial}L(x) = \partial\varphi(x)$. The corresponding variational problem (i.e., minimizing $I(x) = L(x, 0) = \varphi(x) + \varphi^*(0)$) reduces to the classical approach of minimizing a convex functional in order to solve equations of the form $0 \in \partial\varphi(x)$.

More interesting examples of selfdual Lagrangians are of the form $L(x, p) = \varphi(x) + \varphi^*(-\Gamma x + p)$ where φ is a convex and lower semi-continuous function on X , and $\Gamma : X \rightarrow X^*$ is a skew-symmetric operator. The corresponding selfdual vector field is then,

$$\bar{\partial}L(x) = \Gamma x + \partial\varphi(x).$$

More generally, if the operator Γ is merely non-negative (i.e., $\langle \Gamma x, x \rangle \geq 0$), then one can still write the vector field $\Gamma + \partial\varphi$ as $\bar{\partial}M$ for some selfdual Lagrangian M defined now on $X \times X^*$, as

$$M(x, p) = \psi(x) + \psi^*(-\Gamma^{as}x + p)$$

where ψ is the convex function $\psi(x) = \frac{1}{2}\langle \Gamma x, x \rangle + \varphi(x)$, and where $\Gamma^{as} = \frac{1}{2}(\Gamma - \Gamma^*)$ is the anti-symmetric part of Γ , and $\Gamma^{sym} = \frac{1}{2}(\Gamma + \Gamma^*)$ is its symmetric part. The main interest being that equations of the form $\Gamma x + \partial\varphi(x) = p$ can now be solved for a given $p \in X^*$, by simply minimizing the functional

$$I_p(x) = \psi(x) + \psi^*(\Gamma^{as}x + p) - \langle x, p \rangle$$

and proving that its infimum is actually zero.

More classical is the fact that the above examples are typical set-valued nonlinear operators T that are *monotone*, meaning that their graphs $G(T) = \{(x, p) \in X \times X^*; p \in T(x)\}$ are *monotone* subsets of $X \times X^*$, i.e., it satisfies:

$$\langle x - y, p - q \rangle \geq 0 \text{ for every } (x, p) \text{ and } (y, q) \text{ in } G(T). \quad (4)$$

They are actually *maximal monotone operators* meaning that their graph $G(T)$ is maximal in the family of monotone subsets of $X \times X^*$, ordered by set inclusion. For a comprehensive description of this theory, we refer to the monograph of Brezis [1].

It was therefore a natural question whether one can associate a selfdual Lagrangian L to any maximal operator T , so that equations involving such operators can be resolved variationally. We recently realized that Krauss [12] and Fitzpatrick [3] had done some work in this direction in the 80's, and have managed to associate to a maximal monotone operator T , a "sub-selfdual Lagrangian", i.e. a convex lower semi-continuous function L on state space $X \times X^*$ satisfying $L^*(p, x) \geq L(x, p)$ on $X \times X^*$, and in such a way that $T = \bar{\partial}L$. The main interest of this note is to show that one can actually construct a selfdual Lagrangian associated to T , which answers one of the questions in [3].

Theorem 1.1 *Let L be a proper selfdual Lagrangian L on a reflexive Banach space $X \times X^*$, then the vector field $x \rightarrow \bar{\partial}L(x)$ is maximal monotone.*

Conversely, if $T : D(T) \subset X \rightarrow 2^{X^*}$ is a maximal monotone operator with a non-empty domain, then there exists a selfdual Lagrangian N on $X \times X^*$ such that $T = \bar{\partial}N$.

It is worth comparing this result with the following celebrated result of Rockafellar [13], which gives an integral representation of those maximal monotone operators $T : X \rightarrow 2^{X^*}$ that are *cyclically monotone*, i.e., those that verify for any finite number of points $(x_i, p_i)_{i=0}^n$ in the graph $G(T)$ with $x_0 = x_n$, we have

$$\sum_{i=1}^n \langle p_k, x_k - x_{k-1} \rangle \geq 0. \quad (5)$$

Theorem 1.2 (Rockafellar) *If $\varphi : X \rightarrow \mathbb{R} \cup \{+\infty\}$ is a proper convex and lowersemi continuous functional on a Banach space X , then its subdifferential map $x \rightarrow \partial\varphi(x)$ is a maximal cyclically monotone map. Conversely if $T : X \rightarrow 2^{X^*}$ is a maximal cyclically monotone map with a non-empty domain, then there exists a proper convex and lower semi continuous functional on X such that $T = \partial\varphi$.*

The advantages of identifying maximal monotone operators with the class of selfdual vector fields are numerous:

1. As mentioned above, all equations, systems, variational inequalities, and dissipative initial value parabolic problems which traditionally involve maximal monotone operators, can now be formulated and resolved variationally. These problems can therefore be analyzed with the full range of methods –computational or not– that are available for variational settings.
2. While issues around the superposition of, and other operations on, maximal monotone operators are often delicate to prove, the class of selfdual Lagrangians possesses remarkable permanence properties that are also easy to establish. It reflects most variational aspects of convex analysis, and is stable under similar type of operations making the calculus of selfdual Lagrangians as manageable, yet much more encompassing [4].
3. Selfduality allows for the superposition of appropriate boundary Lagrangians with “interior” Lagrangians, leading to the resolution of problems with various linear and nonlinear boundary constraints that are not amenable to the standard variational theory [10]. They can also be iterated with certain nonlinear operators that are far from being maximal monotone [5].
4. Selfdual Lagrangians defined on state spaces also lift to selfdual Lagrangians on path spaces leading to a unified approach for stationary and dynamic equations. More precisely, selfdual flows of the form $\dot{u}(t) \in \bar{\partial}L(u(t))$ with a variety of time-boundary conditions can be reformulated and resolved as $0 \in \bar{\partial}\mathcal{L}(u)$ where $\mathcal{L}$ is a corresponding selfdual Lagrangian on path space, a phenomenon that leads to natural and quite interesting iterations (See [4], [11]).
5. The class of B -selfdual Lagrangians corresponding to a general automorphism B and their B -selfdual vector fields goes beyond the theory of maximal monotone operators. They also cover the theory of Hamiltonian systems –when B is the symplectic matrix J – and other twisted differential operators [7]. In return, the connection between the two concepts leads to a rich parallel theory of B -monotone operators which coincides with the classical theory when B is the identity. The corresponding class of PDEs and their variational principles are studied in detail in the upcoming monograph [6].

2 Selfdual vector fields are maximal monotone operators

To any convex lower semi-continuous Lagrangian $L : X \times X^* \rightarrow \mathbb{R} \cup \{+\infty\}$, we can associate a vector field δL at any $x \in X$, to be the –possibly empty– set

$$\delta L(x) := \{p \in X^*; (p, x) \in \partial L(x, p)\}. \quad (6)$$

It is easy to see that the convexity of L yields that $x \rightarrow \delta L(x)$ is a monotone map, since if $p \in \delta L(x)$ and $q \in \delta L(y)$, then

$$\langle p - q, x - y \rangle = \frac{1}{2} \langle (p, x) - (q, y), (x, p) - (y, q) \rangle \geq 0.$$

We can also associate another –not always necessarily monotone– vector field as:

$$\bar{\partial}L(x) := \{p \in X^*; L(x, p) - \langle x, p \rangle = 0\}. \quad (7)$$

Note that $\bar{\partial}L$ should not be confused with the subdifferential ∂L of L as a convex function on $X \times X^*$. If now L satisfies

$$L(x, p) \geq \langle x, p \rangle \text{ for all } (x, p) \in X \times X^*, \quad (8)$$

then it is easy to see that $\bar{\partial}L(x) \subset \delta L(x)$. Moreover, if L is selfdual, i.e., if

$$L^*(p, x) = L(x, p) \text{ for all } (x, p) \in X \times X^*, \quad (9)$$

then the reverse inclusion is also true since then Legendre-Fenchel duality applied to L yields:

$$2L^*(p, x) = 2L(x, p) = L^*(p, x) + L(x, p) \geq 2\langle x, p \rangle$$

and therefore $L(x, p) - \langle x, p \rangle = 0$ if and only if $L(x, p) + L^*(p, x) = \langle x, p \rangle + \langle p, x \rangle$ which is equivalent to $(p, x) \in \partial L(x, p)$. In other words, we can state the following observation.

Lemma 2.1 *If L is a convex lower semi-continuous selfdual Lagrangian on $X \times X^*$, then $p \in \bar{\partial}L(x)$ if and only if $p \in \delta L(x)$.*

Also note that $p \in \bar{\partial}L(x)$ if and only if $0 \in \bar{\partial}L_p(x)$ where L_p is the selfdual Lagrangian $L_p(x, q) = L(x, p + q) + \langle x, p \rangle$. This is also equivalent to the statement that the infimum of the functional $I_p(u) = L(u, p) - \langle u, p \rangle$ is zero and is attained at $x \in X$. This leads to the following proposition which is a particular case of a more general result established in [4].

Proposition 2.1 *Let L be a convex lower semi-continuous selfdual Lagrangian on a reflexive Banach space $X \times X^*$, such that for some $x_0 \in X$, the function $p \rightarrow L(x_0, p)$ is bounded on the balls of X^* . Then for each $p \in X^*$, there exists $\bar{x} \in X$ such that:*

$$\begin{cases} L(\bar{x}, p) - \langle \bar{x}, p \rangle = \inf_{x \in X} \{L(x, p) - \langle x, p \rangle\} = 0. \\ p \in \bar{\partial}L(\bar{x}). \end{cases} \quad (10)$$

Proof: We can assume that $p = 0$ by considering the translated Lagrangian $M(x, q) = L(x, p + q) - \langle x, p \rangle$ which is also selfdual on $X \times X^*$. In this case $M(x, 0) \geq 0$ for every $x \in X$.

Now consider $(\mathcal{P}_q)$ the primal minimization problem $h(q) = \inf_{x \in X} M(x, q)$ in such a way that $(\mathcal{P}_0)$ is the initial problem $h(0) = \inf_{x \in X} M(x, 0)$, and the dual problem $(\mathcal{P}^*)$ is therefore $\sup_{y \in X} -M^*(0, y)$.

We readily have the following weak duality formula:

$$\inf_{x \in X} \mathcal{P}_0 := \inf_{x \in X} M(x, 0) \geq 0 \geq \sup_{y \in X} -M^*(0, y) := \sup \mathcal{P}^*.$$

Note that h is convex on X^* , and that its Legendre conjugate satisfies for all $y \in X$.

$$h^*(y) = M^*(0, y) = M(y, 0) = L(y, p) - \langle y, p \rangle$$

Moreover, $h(q) = \inf_{x \in X} M(x, q) \leq L(x_0, p + q) - \langle x_0, p \rangle$ and therefore $q \rightarrow h(q)$ is bounded above on the balls of X^* , and hence it is subdifferentiable at 0 (i.e., the problem $(\mathcal{P}_0)$ is then stable). Any point $\bar{x} \in \partial h(0)$ satisfies $h(0) + h^*(\bar{x}) = 0$, which means that

$$-\inf_{x \in X} M(x, 0) = -h(0) = h^*(\bar{x}) = M^*(0, \bar{x}) = M(\bar{x}, 0) \geq \inf_{x \in X} M(x, 0).$$

It follows that $\inf_{x \in X} M(x, 0) = M(\bar{x}, 0) \leq 0$ and the infimum of $(\mathcal{P})$ is therefore zero and is attained at $\bar{x}$.

Lemma 2.2 *Let L be a proper selfdual Lagrangian on $X \times X^*$, then for any convex continuous function φ on X , the Lagrangian defined by*

$$M(x, p) = \inf \{L(x, p - r) + \varphi(x) + \varphi^*(r); r \in X^*\}$$

is also selfdual on $X \times X^*$.

Proof: Indeed, fix $(q, y) \in X^* \times X$ and write:

$$\begin{aligned}
M^*(q, y) &= \sup\{\langle q, x \rangle + \langle y, p \rangle - L(x, p - r) - \varphi(x) - \varphi^*(r); (x, p, r) \in X \times X^* \times X^*\} \\
&= \sup\{\langle q, x \rangle + \langle y, r + s \rangle - L(x, s) - \varphi(x) - \varphi^*(r); (x, s, r) \in X \times X^* \times X^*\} \\
&= \sup_{x \in X} \left\{ \langle x, q \rangle + \sup_{(s, r) \in X^* \times X^*} \{ \langle y, r + s \rangle - L(x, s) - \varphi^*(r) \} - \varphi(x) \right\} \\
&= \sup_{x \in X} \left\{ \langle x, q \rangle + \sup_{s \in X^*} \{ \langle y, s \rangle - L(x, s) \} + \sup_{r \in X^*} \{ \langle y, r \rangle - \varphi^*(r) \} - \varphi(x) \right\} \\
&= \sup_{x \in X} \left\{ \langle x, q \rangle + \sup_{s \in X^*} \{ \langle y, s \rangle - L(x, s) \} + \varphi(y) - \varphi(x) \right\} \\
&= \sup_{x \in X} \sup_{s \in X^*} \{ \langle x, q \rangle + \langle y, s \rangle - L(x, s) - \varphi(x) \} + \varphi(y) \\
&= (L + T_\varphi)^*(q, y) + \varphi(y)
\end{aligned}$$

where $T_\varphi(x, s) := \varphi(x)$ for all $(x, s) \in X \times X^*$. Note now that

$$T_\varphi^*(q, y) = \sup_{x, s} \{ \langle q, x \rangle + \langle y, s \rangle - \varphi(x) \} = \begin{cases} +\infty & \text{if } y \neq 0 \\ \varphi^*(q) & \text{if } y = 0 \end{cases}$$

in such a way that by using the duality between sums and convolutions in both variables, we get

$$\begin{aligned}
(L + T_\varphi)^*(q, y) &= \text{conv}(L^*, T_\varphi^*)(q, y) \\
&= \inf_{r \in X^*, z \in X} \{ L^*(r, z) + T_\varphi^*(-r + q, -z + y) \} \\
&= \inf_{r \in X^*} \{ L^*(r, y) + \varphi^*(-r + q) \}
\end{aligned}$$

and finally

$$\begin{aligned}
M^*(q, y) &= (L_\varphi + T)^*(q, y) + \varphi(y) \\
&= \inf_{r \in X^*} \{ L^*(r, y) + \varphi^*(-r + q) \} + \varphi(y) \\
&= \inf_{s \in X^*} \{ L^*(q - s, y) + \varphi^*(s) \} + \varphi(y) \\
&= \inf_{s \in X^*} \{ L(y, q - s) + \varphi(y) + \varphi^*(s) \} \\
&= M(q, y).
\end{aligned}$$

Proposition 2.2 *Let L be a selfdual Lagrangian L on a reflexive Banach space $X \times X^*$. The following assertions then hold:*

1. *The vector field $x \rightarrow \bar{\partial}L(x)$ is maximal monotone.*
2. *If L is strictly convex in the second variable, then the maximal monotone vector field $x \rightarrow \bar{\partial}L(x)$ is single-valued on its domain.*
3. *If L is uniformly convex in the second variable (i.e., if $L(x, p) - \epsilon \frac{\|p\|^2}{2}$ is convex in p for some $\epsilon > 0$) then the vector field $x \rightarrow \bar{\partial}L(x)$ is a Lipschitz maximal monotone operator on its domain.*

Proof: Denoting by $J : X \rightarrow 2^{X^*}$ the duality map between X and X^* , it suffices to show that the vector field $\bar{\partial}L + J$ is onto [13]. In other words, we need to find for any $p \in X^*$, an $x \in X$ such that $p = \bar{\partial}L(x) + J(x)$. For that, we consider the following Lagrangian on $X \times X^*$.

$$M(x, p) = \inf \left\{ L(x, p - r) + \frac{1}{2} \|x\|_X^2 + \frac{1}{2} \|r\|_{X^*}^2; r \in X^* \right\}.$$

It is a selfdual Lagrangian according to the previous lemma. Moreover – assuming without loss of generality – that the point $(0, 0)$ is in the domain of L , we get the estimate $M(0, p) \leq L(0, 0) + \frac{1}{2}\|p\|_{X^*}^2$, and therefore Proposition 2.1 applies and we obtain $\bar{x} \in X$ so that $p \in \bar{\partial}M(\bar{x})$. This means that

$$M(\bar{x}, p) - \langle \bar{x}, p \rangle = \inf \left\{ L(\bar{x}, p - r) - \langle \bar{x}, p - r \rangle + \frac{1}{2}\|\bar{x}\|_X^2 + \frac{1}{2}\|r\|_{X^*}^2 - \langle \bar{x}, r \rangle; r \in X^* \right\} = 0.$$

In other words, there exists $\bar{r} \in J(\bar{x})$ such that $p - \bar{r} \in \bar{\partial}L(\bar{x})$ and we are done.

The other assertions of the proposition are straightforward and left to the interested reader.

3 Maximal monotone operators are selfdual vector fields

We start with the following lemma which is essentially due to Fitzpatrick [3].

Lemma 3.1 *Let $T : D(T) \subset X \rightarrow 2^{X^*}$ be a monotone operator, and consider on $X \times X^*$ the Lagrangian L_T defined by*

$$L_T(x, p) = \sup \{ \langle p, y \rangle + \langle q, x - y \rangle; (y, p) \in G(T) \} \quad (11)$$

1. *If $\text{Dom}(T) \neq \emptyset$, then L_T is convex and lower semi-continuous function on $X \times X^*$ such that for every $x \in \text{Dom}(T)$, we have $Tx \subset \bar{\partial}L(x) \cap \delta L(x)$. Moreover, we have*

$$L_T^*(p, x) \geq L_T(x, p) \text{ for every } (x, p) \in X \times X^*. \quad (12)$$

2. *If T is maximal monotone, then $T = \bar{\partial}L = \delta L$ and*

$$L_T(x, p) \geq \langle x, p \rangle \text{ for all } (x, p) \in X \times X^*. \quad (13)$$

Proof: (1) If $x \in \text{Dom}(T)$ and $p \in Tx$, then the monotonicity of T yields for any $(y, q) \in G(T)$

$$\langle x, p \rangle \geq \langle y, p \rangle + \langle x - y, q \rangle$$

in such a way that $L_T(x, p) \leq \langle x, p \rangle$. On the other hand, we have

$$L_T(x, p) \geq \langle x, p \rangle + \langle p, x - x \rangle = \langle x, p \rangle,$$

and therefore $p \in \bar{\partial}L(x)$.

Write now for any $(y, q) \in X \times X^*$,

$$\begin{aligned} L_T(x + y, p + q) - L_T(x, p) &= \sup \{ \langle p + q, z \rangle + \langle r, x + y \rangle - \langle z, r \rangle; (z, r) \in G(T) \} - L_T(x, p) \\ &\geq \langle p + q, x \rangle + \langle p, x + y \rangle - \langle p, x \rangle - \langle p, x \rangle \\ &= \langle q, x \rangle + \langle p, y \rangle \end{aligned}$$

which means that $(p, x) \in \partial L(x, p)$ and therefore $p \in \delta L(x)$.

Note also that $L_T(x, p) = H_T^*(p, x)$ where H_T is the Lagrangian on $X \times X^*$ defined by

$$H_T(x, p) = \begin{cases} \langle x, p \rangle & \text{if } (x, p) \in G(T) \\ +\infty & \text{otherwise.} \end{cases} \quad (14)$$

Since $L_T(x, p) = \langle x, p \rangle = H_T(x, p)$ whenever $(x, p) \in G(T)$, it follows that $L_T \leq H_T$ on $X \times X^*$ and so $L_T^*(p, x) \geq H_T^*(p, x) = L_T(x, p)$ everywhere.

- (2) If now T is maximal then necessarily $Tx = \delta L(x) \cap \bar{\partial}L(x) = \delta L(x)$ since $x \rightarrow \delta L(x)$ is a monotone extension of T .

In order to show (13), assume to the contrary that $L_T(x, p) < \langle x, p \rangle$ for some $(x, p) \in X \times X^*$. It follows that

$$\langle p, y \rangle + \langle q, x - y \rangle < \langle p, x \rangle \text{ for all } (y, q) \in G(T),$$

and therefore

$$\langle p - q, x - y \rangle > 0 \text{ for all } (y, q) \in G(T).$$

But since T is maximal monotone, this means that $p \in Tx$. But then $p \in \bar{\partial}L_T(x)$ by the first part, leading to $L_T(x, p) = \langle x, p \rangle$, which is a contradiction.

Finally, note that property (13) on L yields that $\bar{\partial}L(x) \subset \delta L(x)$ and therefore we finally obtain that $Tx = \bar{\partial}L(x)$.

Proposition 3.1 *Let X be a separable reflexive Banach space, and let L be a convex lower semi-continuous Lagrangian on $X \times X^*$ that satisfies*

$$L^*(p, x) \geq L(x, p) \geq \langle x, p \rangle \text{ for every } (x, p) \in X \times X^*. \quad (15)$$

Then, there exists a selfdual Lagrangian N on $X \times X^*$ such that $\bar{\partial}L = \bar{\partial}N$ and

$$L(x, p) \leq N(x, p) \leq L^*(p, x) \text{ for every } (x, p) \in X \times X^*. \quad (16)$$

Proof: Consider the following two Lagrangians $M_0(x, p) = L(x, p)$ and $N_0(x, p) = L^*(p, x)$ in such a way that for every $(x, p) \in X \times X^*$,

$$\langle x, p \rangle \leq M_0(x, p) \leq N_0(x, p), \quad (17)$$

$$M_0^*(p, x) = N_0(x, p), \quad (18)$$

and

$$\bar{\partial}M_0(x) = \bar{\partial}N_0(x). \quad (19)$$

We define by transfinite induction a family of convex lower semi-continuous Lagrangians M_α and N_α that verify the following properties for every ordinal α :

1. $M_0 \leq M_\alpha \leq M_{\alpha+1} \leq N_{\alpha+1} \leq N_\alpha \leq N_0$.
2. $M_\alpha^*(p, x) = N_\alpha(x, p)$.
3. $\bar{\partial}M_\alpha(x) = \bar{\partial}N_\alpha(x)$.

Starting with $M_0(x, p)$ and $N_0(x, p)$, we define $(M_\alpha, N_\alpha)_\alpha$ in the following way:

(i) If $\alpha = \beta + 1$, then set

$$M_\alpha(x, p) = \inf \left\{ \frac{M_\beta(x-y, p-q)}{2} + \frac{N_\beta(x+y, p+q)}{2}; (y, q) \in X \times X^* \right\}$$

and

$$N_\alpha(x, p) = \frac{1}{2}(M_\beta(x, p) + N_\beta(x, p)).$$

It is then clear that

$$M_\beta \leq M_\alpha \leq N_\alpha \leq N_\beta,$$

and that $M_\alpha^*(p, x) = (\frac{M_\beta}{2} \star \frac{N_\beta}{2})^*(2p, 2x) = ((\frac{M_\beta}{2})^* + (\frac{N_\beta}{2})^*)(p, x) = (\frac{N_\beta}{2} + \frac{M_\beta}{2})(x, p) = N_\alpha(x, p)$.

(ii) If α is a limit ordinal, then we define:

$$M_\alpha(x, p) = \sup_{\beta < \alpha} M_\beta(x, p) = \lim_{\beta_n \rightarrow \alpha} \uparrow M_{\beta_n}(x, p) \quad \text{and} \quad N_\alpha(x, p) = \inf_{\beta < \alpha} N_\beta(x, p) = \lim_{\beta_n \rightarrow \alpha} \downarrow N_{\beta_n}(x, p).$$

It is again clear that $M_\beta \leq M_\alpha \leq N_\alpha \leq N_\beta$ for every $\beta < \alpha$, and that $M_\alpha^*(p, x) = N_\alpha(x, p)$. Since X is separable, there exists $\gamma < \Omega$ the first uncountable ordinal such that $M_{\gamma+1} = M_\gamma$, which means that $M_\gamma = N_\gamma$. Set $N(x, p) = M_\gamma(x, p)$ and note that

$$N^*(p, x) = M_\gamma^*(p, x) = N_\gamma(x, p) = M_\gamma(x, p) = N(x, p),$$

which means that N is a selfdual Lagrangian. Moreover, since for every $\alpha < \Omega$ we have

$$N_0(x, p) - \langle x, p \rangle \geq N_\alpha(x, p) - \langle x, p \rangle \geq M_\alpha(x, p) - \langle x, p \rangle \geq M_0(x, p) - \langle x, p \rangle \geq 0,$$

it follows that $\bar{\partial}M_0(x) = \bar{\partial}N_L(x) = \bar{\partial}N_0(x)$.

End of proof of Theorem 1.1: Associate to the maximal monotone operator T the sub-selfdual Lagrangian L_T via Lemma 3.1, that is

$$T = \bar{\partial}L_T \text{ and } L_T^*(p, x) \geq L_T(x, p) \geq \langle x, p \rangle.$$

Now apply the preceding Proposition to L_T to find a selfdual Lagrangian N_T such that $Tx = \bar{\partial}N_T(x)$ for any $x \in \text{Dom}(T)$.

References

- [1] H. Brezis. *Opérateurs maximaux monotones et semi-groupes de contractions dans les espaces de Hilbert*, North Holland, Amsterdam-London, 1973.
- [2] H. Brezis, I. Ekeland, *Un principe variationnel associé à certaines equations paraboliques. Le cas indépendant du temps*, C.R. Acad. Sci. Paris Sér. A **282** (1976), 971–974.
- [3] S.P. Fitzpatrick, *Representing monotone operators by convex functions*, Proc. Centre for Math. Analysis 20 (1989), p. 59–65.
- [4] N. Ghoussoub, *Anti-selfdual Lagrangians: Variational resolutions of non self-adjoint equations and dissipative evolutions*, To appear in AIHP-Analyse non linéaire (2005).
- [5] N. Ghoussoub, *Anti-symmetric Hamiltonians: Variational resolution for Navier-Stokes equations and other nonlinear evolutions*, To appear in Comm. Pure & Applied Math. (2005) 25 pp.
- [6] N. Ghoussoub, *Selfdual partial differential systems and its variational principles*, Monograph (2006) (to appear) 250 pp.
- [7] N. Ghoussoub, A. Moameni, *Selfdual variational principles for periodic solutions of Hamiltonian and other dynamical systems*, To appear in Comm. in PDE (2006)
- [8] N. Ghoussoub, A. Moameni, *Selfduality and periodic solutions of certain Schrödinger equations and infinite dimensional Hamiltonian systems*, submitted (2006)
- [9] N. Ghoussoub, L. Tzou. *A variational principle for gradient flows*, Math. Annalen, Vol 30, 3 (2004) p. 519–549.
- [10] N. Ghoussoub, L. Tzou. *Anti-selfdual Lagrangians II: Unbounded non self-adjoint operators and evolution equations*, Annali di Mat. Pura ed Applicata (2005) 30 pp.
- [11] N. Ghoussoub, L. Tzou. *Iterations of anti-selfdual Lagrangians and applications to Hamiltonian systems and multiparameter gradient flows*, Calc. Var.& PDE Vol. 26, N. 4 (2006) p. 511– 534
- [12] E. Krauss, *A representation of maximal monotone operators by saddle functions*, Rev. Roum. Math. Pures Appl. 309 (1985) p. 823–837
- [13] R.R. Phelps, *Convex functions, monotone operators and differentiability*, Lecture Notes in Math. 1364, Springer Verlag, New York, Berlin, Tokyo, (1998), 2nd edition 1993.